

ANTI-SU(5)

J.-P. Derendinger, Jihn E. Kim*) and D.V. Nanopoulos

CERN — Geneva

ABSTRACT

We discuss ordinary as well as supersymmetric $SU(5)\times\widetilde{U}(1)$ models in the hope of accommodating acceptable τ_p and $\sin^2\theta_W$. The ordinary $SU(5)\times\widetilde{U}(1)$ model does not have the monopole. The supersymmetric $SU(5)\times\widetilde{U}(1)$ model can be unified in SO(10).

^{*)} On leave of absence from the Department of Physics, Seoul National University, Seoul 151, Korea.

1. Grand unified theories (GUTs) provide a well-defined framework capable of unifying weak, electromagnetic and strong interactions $^1)$. The "hard" predictions of GUTs include $\sin^2\theta_W \cong 0.215$, $m_b/m_\tau \sim 2.9$ and proton decay mainly to $e^+\pi^0$ with a lifetime $\tau_p \sim 10^{29\pm 1}$ years. On the other hand, GUTs have a rich topological structure such that superheavy monopoles (M $\sim 10^{16}$ GeV) are contained in the particle spectrum of the theory 2). Present experimental evidence disfavours either proton decay to $e^+\pi^0$ or the existence of monopoles as predicted in the minimal SU(5) $^1), ^4)$.

It is remarkable that by going supersymmetric $^{5)}$, both the above problems are naturally eliminated, while $\sin^2\theta_W$ and m_b/m_τ remain unchanged $^{6)}$. Namely, in SUSY GUTs the p \rightarrow e $^{+}\pi^0$ mode is naturally suppressed (ν K or μ K modes are the favourable channels) while a delayed phase transition from the GUT to the SU(3)×SU(2)×U(1) phase ($T_c \sim 10^{10}$ GeV) or inflation $^{7)}$ evades the monopole problem.

Nevertheless, it is of considerable interest and a challenging problem to construct ordinary GUTs which do not suffer from the above deseases. This is the problem that we address in this paper, out of scientific curiosity, since we are fully aware of the "goodies" of SUSY models. It is by considering the introduction of an extra $\widetilde{U}(1)$ which contains a part of the electromagnetic gauge group $U(1)_{em}$ that there is no stable monopole in the theory and the monopole problem does not exist. Furthermore, the SU(N) coupling constant and the $\widetilde{U}(1)$ coupling constant can be arbitrary and hence the proton lifetime can be made sufficiently longer.

With proper phenomenological inputs, we calculate the SU(N) coupling constant g_N^2 and the $\widetilde{\mathrm{U}}(1)$ coupling constant \widetilde{g}_1^2 at the SU(3)×SU(2) unification scale $\widetilde{\mathrm{M}}$. If $g_N^2 < g_1^2$ the group SU(N)× $\widetilde{\mathrm{U}}(1)$ is the partial unification group, and we achieve our objectives. If $g_N^2 > \widetilde{g}_1^2$, there exists a possibility of further unification of SU(N)× $\widetilde{\mathrm{U}}(1)$. Then we cannot resist unifying it in a simple group at $M_{u} > \widetilde{\mathrm{M}}$, and in this case, the monopole problem is resolved by the inflationary idea 7). Indeed, we encounter both of these examples in SU(5)× $\widetilde{\mathrm{U}}(1)$ models with and without supersymmetry.

The paper is organized as follows. In Section 2, we set our rules for finding fermion representations in $SU(N)\times\widetilde{U}(1)$ models and point out that only one class of models is available for our purpose. In Sections 3 and 4, we present $SU(5)\times\widetilde{U}(1)$ models with and without SUSY, respectively. In Section 5, we show that an $SU(7)\times\widetilde{U}(1)$ model with integer charged leptons is not a viable choice.

- 2. With educated reasons $^{8)-10}$, we set the following rules for $SU(N)\times\widetilde{U}(1)$ theories:
- i) There should not exist triangle anomalies;
- ii) The fermion representation must be chiral under $SU(N)\times \widetilde{U}(1)$,
- iii) The fermion representation must be real under the subgroup $SU(3)_c \times U(1)_{em}$.

Let us concentrate on completely antisymmetric fermion representations of the SU(N) groups. This is reasonable because the quarks are believed to be 3 and 3 * of SU(3) $_{\rm c}$. An irreducible fermion representation with m antisymmetrized indices is denoted as $R_{\rm m}^{\rm N}$. There exist three types of triangle anomalies

$$A_{1}[AAA]_{m}^{N}$$
, $A_{2}[AA\widetilde{Y}]_{m}^{N}$, $A_{3}[\widetilde{Y}\widetilde{Y}\widetilde{Y}]_{m}^{N}$ (1)

for a fermion loop R_m^N where A and \widetilde{Y} inside the bracket represent the external SU(N) gauge bosons or $\widetilde{U}(1)$ gauge bosons. Therefore, we satisfy three anomaly free conditions. From conditions of vanishing A_1 , A_2 and A_3 anomalies, we obtain

$$\sum_{m=1}^{N-1} n_m \frac{(N-2m)(N-3)!}{(N-m-1)!(m-1)!} = 0$$
 (2)

$$\sum_{m=1}^{N-1} n_m \binom{N-2}{m-1} \widetilde{y}_m = 0$$
(3)

$$\sum_{m=1}^{N-1} n_m d_m \tilde{Y}_m^3 + n_o \tilde{Y}_o^3 = 0$$
 (4)

where n_m is the number of irreducible representations R_m^N , and \widetilde{Y}_m is the $\widetilde{U}(1)$ charge of the representation R_m^N . We also introduce an SU(N) singlet R_0^N whose \widetilde{Y} value is \widetilde{Y}_0 . Note that it is generally difficult to satisfy Eqs. (2)-(4) without a singlet. With a singlet(s), Eq. (4) is merely a defining equation for \widetilde{Y}_0 . This definition is possible because a cubic equation has always a real root.

Let us first find out the simplest solution to Eqs. (2)-(4). Equation (4) is satisfied by the introduction of SU(N) singlet(s). The simplest solution is obtained by precise matching of each term in the sum of (2) and (3), which results in the condition

$$\widetilde{Y}_{m} = N - 2m \tag{5}$$

This solution is equivalent to the hypercharges of irreducible representations of SU(N) when a spinor representation of SO(2N) breaks into SU(N)× $\widetilde{\rm U}(1)^{11}$. The simplest choice is therefore $\rm n_m=0$ for $\rm m=odd$ and $\rm n_m=1$ for $\rm m=even$, which is obtainable from one spinor representation of SO(2N). This representation then satisfies the properties ii) and iii) too.

Let us next consider possibilities of more general solutions. For this purpose, the constraints ii) and iii) play important roles. Indeed, there exists a study of this problem in the literature 10 . The conclusion is that (reducible) fermion representations with properties i)-iii) are possible only for the spinor representations of SU(2n+1), i.e., the representations obtained from the spinor representation of SO(4n+2). In Ref. 10), the conclusion was drawn without $\widetilde{\mathrm{U}}(1)$. Nevertheless, we will get the same result with the inclusion of $\widetilde{\mathrm{U}}(1)$ also since the result of Ref. 10) led to the spinor of SU(2n+1). Thus, possible fermion spectra satisfying properties i)-iii) are expected to be:

$$\gamma_{\alpha} + \gamma^{\alpha\beta}$$
 : $SU(5) \times \widetilde{U}(4)$
 $\gamma_{\alpha} + \gamma^{\alpha\beta} + \gamma_{\alpha\beta} \times SU(7) \times \widetilde{U}(4)$ (6)

We show this property explicitly for $SU(5)\times\widetilde{U}(1)$ and $SU(7)\times\widetilde{U}(1)$. The complexity property is apparent from the representations (6). The reality property is equivalent to the existence of all possible Yukawa couplings which can give masses to all $SU(3)_{\rm C}\times U(1)_{\rm em}$ non-trivial fermions. Therefore, we study all possible Yukawa couplings instead of checking the $SU(3)_{\rm C}\times U(1)_{\rm em}$ property. We know that SU(2n+1) fermions are real if all possible Yukawa couplings are allowed. Thus, we prove the reality property by the following dictum. First, write down all possible Yukawa couplings allowed by the SU(2n+1) gauge symmetry only. Then, assign $\widetilde{U}(1)$ hypercharge by the formula (3). If some couplings are forbidden by the $\widetilde{U}(1)$ hypercharge, there is a chance that some fermions do not get masses. If $\widetilde{U}(1)$ hypercharges of Higgs fields are not completely determined, there will remain a global symmetry which forbids mixings between the fermion generations or even some fermions would not get masses. However, if the $\widetilde{U}(1)$ hypercharges of the Higgs fields are uniquely determined, then we obtain the desired reality property.

For $SU(5)\times\widetilde{U}(1)$, the hypercharge assignment by Eq. (3) is identical to the one by Eq. (5), i.e.,

$$\widetilde{Y}(Y_{\alpha}) = -3 , \ \widetilde{Y}(Y^{\alpha}) = 1$$

The Yukawa couplings are

which uniquely determine

$$\widetilde{Y}(H^{\alpha}) = -2 \tag{9}$$

Therefore, we satisfy the reality condition.

For $SU(7)\times \widetilde{U}(1)$, we can satisfy Eq. (3) by the following assignment

$$\widetilde{Y}(Y_{X/3}Y) = -1$$

$$\widetilde{Y}(Y_{X/3}Y) = Y$$

$$\widetilde{Y}(Y_{X/3}Y) = 10 - 5y$$
(10a,b,c)

The relevant Yukawa couplings are

Equations (10) and (11) are satisfied with y = 3, i.e.,

$$\widetilde{Y}(\gamma_{x}) = -5 , \quad \widetilde{Y}(\gamma^{\alpha\beta}) = 3 , \quad \widetilde{Y}(\gamma_{\alpha\beta}) = -1$$

$$\widetilde{Y}(H^{\alpha}) = -2 , \quad \widetilde{Y}(H^{\alpha\beta}) = -6$$

which agrees with the assignment (5). The spinor representation is real.

We have also checked this reality property for $SU(9) \times \widetilde{U}(1)$. It is believed that the spinor representation of $SU(2n+1)\times \widetilde{U}(1)$ with the hypercharge given by Eq. (5) satisfies properties i)-iii) provided a SU(2n+1) singlet has $\widetilde{Y}(\psi_0) = 2n+1.$

Let us consider an $SU(5)\times\widetilde{U}(1)$ model *) without SUSY. This model has the same particle assignment as Barr's 13), but we differ in philosophy from his by not unifying $\widetilde{\mathrm{U}}(1)$ within $\mathrm{SO}(10)$. The electromagnetic charge Q_{em} is given by

 $^{^{*)}}$ In another context, SU(5)×U(1) was considered in Ref. 12).

$$Q_{em} = I_3 - \frac{1}{5}Y' + \frac{1}{5}\widehat{Y}$$
 (13)

where

1:
$$Y' = 0$$
, $\tilde{Y} = 5$ (14a,b)

$$\overline{5}$$
: $Y' = (1/3, 1/3, 1/3; -1/2, -1/2)$ (15a)

$$\tilde{Y} = (-3, -3, -3, -3)$$
 (15b)

10:
$$Y' = \left(-\frac{2}{3}, -\frac{2}{3}, -\frac{2}{3}, \frac{1}{6}, \frac{1}{$$

$$\hat{Y} = (1, 1, 1; 1, 1, 1, 1, 1; 1)$$
 (16b)

Defining the coupling constants associated with T $_3$, Y' and \widetilde{Y} by g_2 , g' and \widetilde{g} , respectively, we have a relation 14)

$$\frac{1}{e^2} = \frac{1}{g_2^2} + \frac{1/25}{g'^2} + \frac{1/25}{\tilde{g}^2}$$
 (17)

To study and compare the evolution of coupling constants, it is useful to define properly normalized generators $Y_1' = C'Y'$ and $\widetilde{Y}_1 = \widetilde{CY}$ such that (on the sixteen states)

$$Tr(Y_1'^2) = Tr(\widehat{Y}_1^2) = Tr(I_3^2) = 2$$
 (18)

(i.e., $C^{2} = 3/5$ and $\tilde{C}^{2} = 1/40$) and the associated coupling constants verify

$$\frac{1}{e^2} = \frac{1}{g_2^2} + \frac{1/15}{g_1^{'2}} + \frac{8/5}{\tilde{g}_1^{'2}}$$
 (19)

In particular, we obtain for $\sin^2\theta_W$ at the SU(5) unification scale \widetilde{M} where $g_2(\widetilde{M})=g_1^*(\widetilde{M})\equiv g_5$

$$\sin^2 \theta_{\rm W}^{\circ} = \frac{3}{8} \cdot \frac{1}{1 + \frac{3}{5} \left[\left(\frac{95}{3} \right)_{\rm Pl}^2 - 1 \right]}$$
(20)

where g_5 is the unification coupling constant at \widetilde{M} . If $\widetilde{g}_1 = g_5$, $\sin^2\theta_W^0 = 3/8$ as expected. To have a larger proton decay rate than the one of the SU(5) model, we must start with a relation $\left|g_5\right| < \left|\widetilde{g}_1\right|$ at \widetilde{M} so that the prediction of $\sin^2\theta_W(M_W)$ is untouched with a larger GUT gap, $M_W-\widetilde{M}$. Because of the condition $\left|g_5\right| < \left|\widetilde{g}_1\right|$, we cannot further unify SU(5)× \widetilde{U} (1).

The evolution of coupling constants is

$$\frac{1}{g_3^2(M_w)} = \frac{1}{g_5^2} + \frac{1}{8\pi^2} \left[-11 + \frac{4}{3} N_g \right] \ln \left(\frac{\widetilde{M}}{M_w} \right)$$
(21)

$$\frac{1}{g_2^2(M_W)} = \frac{1}{g_5^2} + \frac{1}{8\pi^2} \left(-\frac{22}{3} + \frac{4}{3}N_g + \frac{N_H}{6} \right) l_H \left(\frac{\tilde{H}}{M_W} \right)^{(22)}$$

$$\frac{1}{g_1^2(M_W)} = \frac{1}{g_1^2(\widetilde{M})} + \frac{1}{8\pi^2} \left(\frac{4}{3}N_g + \frac{1}{10}N_H\right) \ln\left(\frac{\widetilde{M}}{M_W}\right)^{(23)}$$

where N and N are numbers of families and Higgs doublets. The value $g_1^2(\widetilde{M})$ is related to g_5^2 and $\widetilde{g}_1^2(\widetilde{M})$ by

$$\frac{1}{g_1^2(\widehat{H})} = \frac{1/25}{g_5^2(\widehat{M})} + \frac{24/25}{\widehat{g}_1^2(\widehat{M})}$$
(24)

From (21) and (22), we obtain a useful relation

$$\ln\left(\frac{\widetilde{M}}{M_W}\right) = \frac{2TT\left(\sin^2\theta_W/\alpha_{em} - \frac{1}{4}\alpha_c\right)}{11/3 + NH/6} M_W$$
(25)

For various input parameters of N_H, $\sin^2\theta_W(M_W)$ and $\alpha_c(M_W)$, we present in Table 1 the values \widetilde{M} , $\sin^2\theta_W(\widetilde{M})$, and $(g_5^2/\widetilde{g}_1^2)_{\widetilde{M}}$. For example, for N_g = 3, N_H = 1, $\sin^2\theta_W(M_W)$ = 0.215, $\alpha_c(M_W)$ = 0.13 and $\alpha_{em}(M_W)^{-1}$ = 128, we obtain $\widetilde{M} \cong 9 \times 10^{15}$ GeV, $\tau_p \approx 10^{35}$ years, and $(g_5^2/\widetilde{g}_1^2)_{\widetilde{M}} = 0.97$. If higher order effect does not change the relation $(g_5^2/\widetilde{g}_1^2)_{\widetilde{M}} < 1$, we cannot unify SU(5)× \widetilde{U} (1) in a simple group for this attractive set of input parameters *. In this case, there would not exist a monopole and τ_p is too long to be observed by current proton decay detectors.

4. For the case of SUSY SU(5) $\times \widetilde{U}(1)$, we obtain

$$\frac{1}{g_3^2(M_W)} = \frac{1}{g_5^2} + \frac{1}{8\Pi^2} \left(-9 + 2N_g\right) \ln\left(\frac{M}{M_W}\right)$$
(26)

$$\frac{1}{g_2^2(M_W)} = \frac{1}{g_5^2} + \frac{1}{8\pi^2} \left(-6 + 2N_g + \frac{1}{2}N_H \right) l_H \left(\frac{\widetilde{H}}{H_W} \right)$$
(27)

$$\frac{1}{g_{1}^{2}(M_{W})} = \frac{1}{g_{1}^{2}(\widetilde{H})} + \frac{1}{8\pi^{2}} \left(2N_{g} + \frac{3}{10}N_{H}\right) \ln\left(\frac{\widetilde{M}}{M_{W}}\right)$$
(28)

$$\ln\left(\frac{\widetilde{M}}{Mw}\right) = \frac{2\pi(\sin^2\theta_w/\alpha_{em} - 1/\alpha_e)}{3 + N_H/2}$$
(29)

^{*)}However, note that we can have $(g_5^2/\tilde{g}_1^2)_{\widetilde{M}} > 1$ for $N_H = 2$.

In Table 2, we present the values of \widetilde{M} , $\sin^2\theta_W(\widetilde{M})$ and $(g_5^2/\widetilde{g}_1^2)_{\widetilde{M}}$ for several input parameter sets. We note that for the case of $N_H = 2$ reasonable values of \widetilde{M} are obtained from acceptable values of $\sin^2\theta_W(M_W)$. Furthermore, it generally gives $(g_5^2/\widetilde{g}_1^2)_{\widetilde{M}} > 1$, implying a possibility of unification in SO(10). For example, for $N_g = 3$, $N_H = 2$, $\sin^2\theta_W(M_W) = 0.215$, $\alpha_c(M_W) = 0.13$, and $\alpha_{em}(M_W)^{-1} = 128$, we obtain $\widetilde{M} \cong 2.5 \times 10^{15}$ GeV, $\tau_p \cong 10^{33}$ years, and $(g_5^2/\widetilde{g}_1^2)_{\widetilde{M}} \cong 1.32$. The case $N_H = 4$ is not successful.

5. In this section, we present an analysis for the $SU(7)\times\widetilde{U}(1)$ model based on the fermion spectrum of Ref. 15). For the fermion spectrum obtainable from SO(4n+2), it is generally true to have only two patterns of $SU(2n+1)\times\widetilde{U}(1)$: one is the usual SU(2n+1) and the other is the anti- $SU(2n+1)\times\widetilde{U}(1)$. The Dynkin weight diagram of the spinor of SO(4n+2) has a distinctive shape SU(2n+1). The interconnected central part is connected to two strings with two weights on each string. The weight on the end of a string is either the highest or the lowest weight. Only these two weights can be singlets under $SU(2n+1)\times\widetilde{U}(1)$, since we can disconnect only one simple root from either of these two to make the root a singlet under $SU(2n+1)\times\widetilde{U}(1)$. The analysis presented in this part can be applied to $SU(7)\times\widetilde{U}(1)$ models with fractionally charged leptons.

The anti- $SU(7)\times\widetilde{U}(1)$ model has the following relations,

$$Q_{em} = I_3 + Y' + \frac{1}{7} \widetilde{Y}$$
 (30)

$$Y'(7) = diag(\frac{13}{21}, \frac{13}{21}, \frac{13}{21}; \frac{3}{14}; \frac{3}{14}; \frac{5}{7}, \frac{5}{7})$$
(31)

$$\widetilde{Y}(1) = 7$$
 , $\widetilde{Y}(\overline{7}) = -5$
 $\widetilde{Y}(21) = 3$, $\widetilde{Y}(\overline{35}) = -1$ (32)-(35)

$$\frac{1}{e^2} = \frac{1}{g_2^2} + \frac{\frac{1}{8} \sum Y'^2}{g_1'^2} + \frac{\frac{1}{49} \frac{1}{8} \sum \widetilde{Y}^2}{\widetilde{g}_1^2}$$
(36)

where $\mathbf{g_1'}$ and $\overset{\sim}{\mathbf{g}_1}$ are the coupling constants for properly normalized generators. Since

$$Tr Y'^{2}(4\alpha) = \frac{95}{42}$$
 (37)

Tr
$$Y'^{2}(Y^{\alpha\beta}) = (N-2) \text{Tr } Y'^{2}(Y^{\alpha})$$
 (38)

Tr
$$Y'^{2}(\gamma^{\alpha\beta\delta}) = \frac{(N-2)(N-3)}{2} \text{Tr } Y'^{2}(\gamma^{\alpha})$$
 (39)

we have

Tr
$$Y'^{2}(64) = 16.\frac{95}{42}$$
 (40)

$$Tr \ \widetilde{Y}^{2} (64) = 16.28$$
 (41)

Therefore,

$$\frac{1}{e^2} = \frac{1}{g_2^2} + \frac{95/21}{g_1'^2} + \frac{8/7}{\tilde{g}_1^2}$$
(42)

$$\sin^2 \theta_W^\circ = \frac{21/116}{1 + \frac{6}{29} (97^2/3^2) \tilde{M}}$$
 (43)

Certainly, we obtain $\sin^2 \theta_W^0 = 3/20$ for $g_7 = \widetilde{g}_1$.

The renormalization group analysis of coupling constants does not give acceptable intermediate mass scales, and we do not succeed in the $SU(7)\times\widetilde{U}(1)$ model. However, it will be certainly possible to have acceptable intermediate scales for $SU(7)\times\widetilde{U}(1)$ models with fractionally charged leptons.

6. We have seen that adding supersymmetry to the $SU(5)\times\widetilde{U}(1)$ model leaves open the possibility of subsequent unification. At \widetilde{M} , $g_5/\widetilde{g}_1>1$, and then these coupling constants will meet at some new scale $M_u>\widetilde{M}$, where unification into SO(10) for instance can occur. We will now discuss the supersymmetric SO(10) model and the possible realization of the symmetry breaking pattern

$$SO(10) \longrightarrow SU(5) \times \widetilde{U}(1)$$

$$\longrightarrow SU(3)_{c} \times SU(2)_{L} \times U(1)_{y}$$
(44)

SO(10) breaking into SU(5)× $\widetilde{\rm U}(1)$ can, in principle, be induced by a real antisymmetric tensorial representation with an even number of indices, i.e., $\frac{45}{45}$ or $\frac{210}{210}$, denoted generically by ϕ . $\frac{45}{45}$ does not possess a cubic invariant. Thus, the corresponding superpotential is only a mass term $\frac{1}{2}$ M Tr ϕ^2 , which leads to the vanishing vacuum expectation value (v.e.v.). This negative result can be corrected by coupling $\frac{45}{45}$ to other multiplets, price to be paid being that the scale and the little group of the v.e.v. is fixed by this new sector of the model. This problem does not occur with $\frac{210}{210}$ where the superpotential reads $\frac{1}{2}$ M Tr ϕ^2 + $\lambda/3$ Tr ϕ^3 leading to a v.e.v. of order M/λ which breaks, among other possibilities, SO(10) into SU(5)× $\widetilde{\rm U}(1)$.

For the second step of symmetry breaking, $SU(5)\times \widetilde{U}(1)\to SU(3)\times SU(2)\times U(1)$, the two natural Higgs candidates are 16 and 126, the desired v.e.v. being in the 10_1 part of 16, and in 50_2 of 126. However, the 126 is more attractive, since its coupling to quark and lepton supermultiplets will give a large Majorana mass to right-handed neutrinos. This is not the case using 16, and since the non-renormalization properties of SUSY suppress radiative corrections, one would get an unacceptable spectrum for neutrinos. Then, using 210 and $126+\overline{126}$ (denoted respectively by ϕ , ψ , ψ), the most general cubic (i.e., renormalizable) superpotential is

$$W = \frac{M}{2} Tr \phi^2 + \frac{\lambda}{3} Tr \phi^3 + \sqrt{T} \phi^4 + \sqrt{T} \phi^4 + \sqrt{T} \phi^4$$
(45)

^{*)} Supersymetric SO(10) models with different patterns of symmetry breaking have been considered in Refs. 16) and 17).

with indices and gamma matrices omitted for clarity. We need both 126 and 126 chiral multiplets to have a superpotential for these fields and to cancel the v.e.v. of the gauge part of the potential 17. This latter requirement enforces the v.e.v.s of 126 and 126 to have the same scale and the same little group. Solving the minimum equations 18, $(\partial W/\partial \phi) = (\partial W/\partial \phi) = 0$ for this superpotential, leads to two difficulties. First, the natural solution is to obtain unbroken SU(5). The $\overline{\phi}\phi\phi$ coupling has in fact a tendency to align the v.e.v.s of 210 (or 45) and 126. The second problem is that this superpotential does not possess two actual scales. The v.e.v. of ϕ is of order M/λ . However, $(\partial W/\partial \phi) = 0$ leads to $<\phi>\approx \mu/\alpha$ and a tuning of parameters is necessary. To solve these problems, we need non-renormalizable terms like, for instance,

$$\frac{\mathcal{S}}{M_{\mathbf{p}}} \overline{\mathbf{T}} \mathbf{Y} \overline{\mathbf{Y}} \mathbf{Y}$$
(46)

To obtain $<\phi>\approx M/\lambda=0(10^{17}~\text{GeV})$ and $<\phi>=0(10^{16}~\text{GeV})$, we will have to impose $\alpha/\beta<0(10^{-4})$ and $\mu/\alpha>0(10^{17}~\text{GeV})$. Notice that choosing $\alpha=0$ leads to pseudo-Goldstone multiplets. Such non-renormalizable terms are naturally obtained in supergravity unified models 6).

5. CONCLUSIONS

The anti-SU(5) models, the ordinary and supersymmetric ones, can be realistic unified models with acceptable $\sin^2\theta_W$ and τ_p . The ordinary SU(5)× $\widetilde{\rm U}(1)$ model is not unified in a simple group and hence there is no stable monopole. The supersymmetric SU(5)× $\widetilde{\rm U}(1)$ model can be realistically unified in the SO(10) group. Other SU(N)× $\widetilde{\rm U}(1)$ models without fractionally charged leptons cannot be made realistic.

 $\frac{\text{TABLE 1}}{\text{M and } (g_5^2/\widetilde{g}_1^2)_{\widetilde{M}}} \text{ in ordinary SU(5)} \times \widetilde{\text{U}(1)}.$

| N _H | $\sin^2\theta_{ m W}({ m M}_{ m W})$ | α _c (M _W) | М | $\sin^2\theta_{\widetilde{W}}(\widetilde{M})$ | $(g_5^2/\widetilde{g}_1^2)_{\widetilde{M}}$ |
|----------------|--------------------------------------|----------------------------------|-----------------------|---|---|
| 1 | 0.215 | 0.10 | 2.18×10 ¹⁴ | 0.359 | 1.074 |
| 1 | 0.215 | 0.13 | 9.58×10 ¹⁵ | 0.382 | 0.970 |
| 1 | 0.215 | 0.16 | 1.02×10 ¹⁷ | 0.397 | 0.910 |
| 1 | 0.225 | 0.10 | 1.73×10 ¹⁵ | 0.382 | 0.969 |
| 1 | 0.225 | 0.13 | 7.59×10 ¹⁶ | 0.405 | 0.875 |
| 1 | 0.225 | 0.16 | 8.07×10 ¹⁷ | 0.420 | 0.820 |
| 2 | 0.215 | 0.10 | 6.63×10 ¹³ | 0.351 | 1.116 |
| 2 | 0.215 | 0.13 | 2.49×10 ¹⁵ | 0.372 | 1.012 |
| 2 | 0.215 | 0.16 | 2.40×10 ¹⁶ | 0.386 | 0.952 |
| 2 | 0.225 | 0.10 | 4.81×10 ¹⁴ | 0.373 | 1.009 |
| 2 | 0.225 | 0.13 | 1.81×10 ¹⁶ | 0.395 | 0.914 |
| 2 | 0.225 | 0.16 | 1.74×10 ¹⁷ | 0.410 | 0.859 |
| <u> </u> | | | | | |

 $\frac{\text{TABLE 2}}{\text{M} \text{ and } (g_5^2/\widetilde{g}_1^2)_{\widetilde{M}} \text{ in SUSY SU(5)} \times \widetilde{\text{U}(1)}}.$

| N _H | $\sin^2\theta_{\mathrm{W}}(\mathrm{M}_{\mathrm{W}})$ | $\alpha_{c}(M_{W})$ | M | $\sin^2 \theta_{W}(M)$ | $(g_5^2/\tilde{g}_1^2)_{\tilde{M}}$ |
|----------------|--|---------------------|-----------------------|------------------------|-------------------------------------|
| 2 | 0.215 | 0.10 | 6.63×10 ¹³ | 0.294 | 1.456 |
| 2 | 0.215 | 0.13 | 2.49×10 ¹⁵ | 0.315 | 1.316 |
| 2 | 0.215 | 0.16 | 2.40×10 ¹⁶ | 0.330 | 1.225 |
| 2 | 0.225 | 0.10 | 4.81×10 ¹⁴ | 0.322 | 1.275 |
| 2 | 0.225 | 0.13 | 1.81×10 ¹⁶ | 0.347 | 1.136 |
| 2 | 0.225 | 0.16 | 1.74×10 ¹⁷ | 0.365 | 1.046 |
| 4 | 0.215 | 0.10 | 2.76×10 ¹¹ | 0.251 | 1.823 |
| 4 | 0.215 | 0.13 | 5.02×10 ¹² | 0.261 | 1.730 |
| 4 | 0.215 | 0.16 | 3.07×10 ¹³ | 0.268 | 1.667 |
| 4 | 0.225 | 0.10 | 1.34×10 ¹² | 0.272 | 1.631 |
| 4 | 0.225 | 0.13 | 2.44×10 ¹³ | 0.284 | 1.533 |
| 4 | 0.225 | 0.16 | 1.49×10 ¹⁴ | 0.293 | 1.467 |
| | | | | | |

REFERENCES

- H. Georgi and S.L. Glashow, Phys. Rev. Lett. 32 (1974) 438;
 H. Georgi, H. Quinn and S. Weinberg, Phys. Rev. Lett. 33 (1974) 450;
 A. Buras, J. Ellis, M.K. Gaillard and D.V. Nanopoulos, Nucl. Phys. B135 (1978) 66;
 J.C. Pati and A. Salam, Phys. Rev. D8 (1973) 1240: D10 (1974) 275.
- 2) G. 't Hooft, Nucl. Phys. B79 (1974) 276; A.M. Polyakov, JETP Lett. 20 (1974) 194.
- 3) R.M. Bionta et al., Phys. Rev. Lett. 51 (1983) 27.
- 4) For a review, see:
 - P. Langacker, Phys. Rep. 72 (1981) 185;
 - D.V. Nanopoulos, Ecole d'Eté de Physique des Particules, Gif-sur-Yvette, 1980 (IN2P3, Paris, 1980), p. 1;
 - J. Ellis, in "Gauge Theories and Experiments at High Energies", eds. K.C. Bowler and D.G. Sutherland, (Scottish Univ. Summer School in Physics, Edinburgh, 1981), p. 201.
- 5) S. Dimopoulos and H. Georgi, Nucl. Phys. B193 (1981) 150;
 N. Sakai, Z. Phys. C11 (1982) 153;
 E. Witten, Nucl. Phys. B185 (1981) 513.
- 6) For recent reviews, see:
 R. Barbieri and S. Ferrara, CERN preprint TH.3547 (1983);
 D.V. Nanopoulos, CERN preprint TH.3699 (1983);
 J. Ellis, CERN preprint TH.3718 (1983).
- 7) A. Guth, Phys. Rev. D23 (1981) 347;
 A.D. Linde, Phys. Lett. 108B (1982) 389;
 A. Albrecht and P.J. Steinhardt, Phys. Rev. Lett. 48 (1982) 1220;
 J. Ellis, D.V. Nanopoulos, K.A. Olive and K. Tamvakis, Nucl. Phys. B221 (1983) 524;
 D.V. Nanopoulos, K.A. Olive and M. Srednicki, Phys. Lett. 127B (1983) 30.
- 8) S.L. Adler, Phys. Rev. 177 (1969) 2426: J.S. Bell and R. Jackiw, Nuovo Cim. 60A (1969) 47.
- 9) H. Georgi, Nucl. Phys. B156 (1979) 126.
- 10) J.E. Kim, J. Kim, K.S. Soh and H.S. Song, Nucl. Phys. B181 (1981) 531.
- 11) R. Slansky, Phys. Rep. 79 (1981) 1.
- 12) A. Zee and J.E. Kim, Phys. Rev. D21 (1980) 1939.
- 13) A. De Rujula, H. Georgi and S.L. Glashow, Phys. Rev. Lett. 45 (1980) 413; S.M. Barr, Phys. Lett. 112B (1982) 219.
- 14) J.E. Kim and H.S. Song, Phys. Rev. D22 (1980) 1753.
- 15) J.E. Kim, Phys. Rev. Lett. 45 (1980) 1916; Phys. Rev. D23 (1981) 2706; D26 (1982) 674.

- 16) L. Ibañez, Phys. Lett. 114B (1982) 243;
 - T.E. Clark, T.K. Kuo and N. Nakagawa, Phys. Lett. 115B (1982) 26;
 - C.S. Aulakh and R.N. Mohapatra, Phys. Rev. D28 (1983) 217;
 - B. Sathiapalan, Caltech preprint 68-922 (1982);
 - J. Maalampi and J. Pulido, to appear in Nucl. Phys. B.
- 17) F. Buccella, J.-P. Derendinger, S. Ferrara and C.A. Savoy, "Unification of the Fundamental Particle Interactions II", eds. J. Ellis and S. Ferrara, (Plenum Press, New York and London, 1983), p. 349; Phys. Lett. 115B (1982) 375;
 - P.H. Frampton and T. Kephart, Phys. Rev. Lett. 48 (1982) 1237; Nucl. Phys. B211 (1983) 239.
- 18) G. Anastaze and J.-P. Derendinger, unpublished.